# Your Guide to Understanding Genetic Conditions

# MBL2 gene

mannose binding lectin 2

#### **Normal Function**

The *MBL2* gene provides instructions for making a protein that assembles into a protein complex called mannose-binding lectin. Functional mannose-binding lectins are made up of two to six protein groups called trimers, which are each composed of three of the protein pieces (subunits) produced from the *MBL2* gene. This protein complex plays an important role in the immune system's response to foreign invaders (pathogens).

Mannose-binding lectin recognizes and attaches (binds) to sugars, such as mannose, fucose, and glucose, that are found on the surface of bacteria, viruses, and yeast. This binding turns on (activates) the complement system, which is a group of immune system proteins that work together to destroy pathogens, trigger inflammation, and remove debris from cells and tissues. Attachment of mannose-binding lectin also targets the pathogen to be engulfed and broken down by special immune cells. Recognition of foreign invaders by mannose-binding lectin provides one of the body's first lines of defense against infection.

### **Health Conditions Related to Genetic Changes**

mannose-binding lectin deficiency

Several common mutations of the *MBL2* gene can lead to a condition called mannose-binding lectin deficiency. People with this condition have low levels of mannose-binding lectin and may be susceptible to recurrent infections. Several of the disease-associated mutations occur in a region of the *MBL2* gene known as exon 1 and result in a change to single protein building blocks (amino acids) in the mannose-binding lectin subunit. Other mutations occur in an area of DNA near the *MBL2* gene called the promoter region, which helps control the production of the mannose-binding lectin subunit.


The change of a single amino acid in the mannose-binding lectin subunit eliminates its ability to assemble into the functional mannose-binding lectin. Similarly, certain mutations in the promoter region of the *MBL2* gene reduce production of the mannose-binding lectin subunit, leading to a decreased number of subunits available for protein assembly and a reduction in the amount of functional protein. With decreased levels of mannose-binding lectin, the body does not recognize and fight foreign invaders efficiently. Consequently, infections can be more common in people with this condition. However, researchers believe that a number of factors,

including other genetic and environmental factors, are involved in the development of mannose-binding lectin deficiency.

#### **Chromosomal Location**

Cytogenetic Location: 10q21.1, which is the long (q) arm of chromosome 10 at position 21.1

Molecular Location: base pairs 52,764,977 to 52,772,845 on chromosome 10 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

#### Other Names for This Gene

- COLEC1
- collectin-1
- HSMBPC
- mannan-binding lectin
- mannose-binding lectin (protein C) 2, soluble
- mannose-binding lectin (protein C) 2, soluble (opsonic defect)
- mannose-binding lectin 2, soluble (opsonic defect)
- mannose-binding protein C
- mannose-binding protein C precursor
- MBL
- MBL2 HUMAN
- MBL2D
- MBP
- MBP-C
- MBP1

#### **Additional Information & Resources**

#### **Educational Resources**

- Immunobiology: The Immune System in Health and Disease (5th Edition, 2001): Mannan-Binding Lectin (MBL) Binds to Patterns of Carbohydrate Groups in the Correct Spatial Orientation https://www.ncbi.nlm.nih.gov/books/NBK27129/figure/A196/
- Molecular Biology of the Cell (4th Edition, 2002): Complement Activation Targets Pathogens for Phagocytosis or Lysis https://www.ncbi.nlm.nih.gov/books/NBK26846/#A4679

#### Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28MBL2%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D

#### **OMIM**

 LECTIN, MANNOSE-BINDING, SOLUBLE, 2 http://omim.org/entry/154545

#### Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology http://atlasgeneticsoncology.org/Genes/GC\_MBL2.html
- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=MBL2%5Bgene%5D
- HGNC Gene Family: Collectins http://www.genenames.org/cgi-bin/genefamilies/set/491
- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene\_symbol\_report?q=data/ hqnc data.php&hqnc id=6922
- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/4153
- UniProt http://www.uniprot.org/uniprot/P11226

#### **Sources for This Summary**

- Arora M, Munoz E, Tenner AJ. Identification of a site on mannan-binding lectin critical for enhancement of phagocytosis. J Biol Chem. 2001 Nov 16;276(46):43087-94. Epub 2001 Aug 30. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/11533031
- Bouwman LH, Roep BO, Roos A. Mannose-binding lectin: clinical implications for infection, transplantation, and autoimmunity. Hum Immunol. 2006 Apr-May;67(4-5):247-56. Epub 2006 Apr 17. Review.
  - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16720204
- OMIM: LECTIN, MANNOSE-BINDING, SOLUBLE, 2 http://omim.org/entry/154545
- Martin P, Lerner A, Johnson L, Lerner DL, Haraguchi S, Good RA, Day NK. Inherited mannose-binding lectin deficiency as evidenced by genetic and immunologic analyses: association with severe recurrent infections. Ann Allergy Asthma Immunol. 2003 Oct;91(4):386-92.
  Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/14582818
- Weis WI, Drickamer K, Hendrickson WA. Structure of a C-type mannose-binding protein complexed with an oligosaccharide. Nature. 1992 Nov 12;360(6400):127-34.
   Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/1436090

## Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/gene/MBL2

Reviewed: March 2012 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services